

## Analysis of Grains

<b>PROTEIN ROUGHAGES</b>	<b>PROTEIN</b>	<b>RUMMINANTS ENERGY mj/kg</b>	<b>POULTRY A.M.E. mj/kg</b>	<b>PIGS A.D.E. mj/kg</b>	<b>FAT</b>	<b>FIBRE</b>	<b>ASH</b>
--------------------------	----------------	------------------------------------	---------------------------------	------------------------------	------------	--------------	------------

### SILAGE

Oaten Hay	5.8	7.5	-	-	2.6	27.0	6.7
Wheaten Hay	6.0	6.7	-	-	2.3	28.0	6.0
Pasture Hay	6.0	7.0	-	-	1.3	29.0	6.1
Oat, Barley or Wheat Straw	3.0	5.0	-	-	1.9	33.9	4.9
Sorghum Stubble	3.6	4.5	-	-	1.3	31.1	11.8
Cottonseed Hulls	4.2	4.8	-	-	0.9	48.0	2.9
Cottonseed	19.0	14.5	-	-	19.6	16.7	3.9
Corn Stubble	4.8	4.5	-	-	1.3	31.1	11.8
Soybean Stubble	5.5	4.5	-	-	1.3	31.1	11.8
Peanut Hulls	3.3	1.3	-	-	3.3	57.2	3.8
Oat Hulls	3.8	5.0	-	-	1.0	33.0	4.0
Peanut Hay	9.3	7.1	-	-	3.3	30.3	8.2
Soybean Hay (Mature)	8.1	5.0	-	-	1.3	38.9	5.6
Wheat/Barley/Rice/Oat Stubble	-	4.5	-	-	1.3	31.1	11.8
Lucerne Hay	15.0	6.9	2.9	5.5	1.6	26.1	9.3
Clover Hay	13.0	8.0	-	-	3.3	22.6	7.1
Soybean Hay (Full Pods)	13.5	7.9	-	-	2.3	30.6	7.2

<b>MISCELLANEOUS</b>	<b>PROTEIN</b>	<b>RUMMINANTS ENERGY mj/kg</b>	<b>POULTRY A.M.E. mj/kg</b>	<b>PIGS A.D.E. mj/kg</b>	<b>FAT</b>	<b>FIBRE</b>	<b>ASH</b>
Molasses	3.5	8.9	9.1	10.3	-	-	10.0
Poultry Litter	25.0	-	-	-	1.3	26.0	16.0

<b>FURNEYS STOCKFEEDS</b>	<b>PROTEIN</b>	<b>RUMMINANTS ENERGY mj/kg</b>	<b>POULTRY A.M.E. mj/kg</b>	<b>PIGS A.D.E. mj/kg</b>	<b>FAT</b>	<b>FIBRE</b>	<b>ASH</b>
Sheep & Cattle Nuts	14.0	12.5	-	-	4	9	-
Lamb Fattener/Cattle Weaner	16.0	12.0	-	-	4	9	-

A.M.E.— Apparent Metabolisable Energy

A.D.E.— Apparent Digestible Energy

Expressed on an “As Fed” basis.

NB:- The above figures are only a guide, protein and energy levels will vary from crop to crop and with different varieties.