

# The Brew 8%

## Dry Season Loose Lick

**A loose supplement that contains bypass protein for weaners in the 100 to 200kg range and sheep of all ages. The Brew contains a small amount of urea to improve rumen development. The Brew assists in supplying the necessary protein, energy and minerals to ensure your sheep and cattle have better growth in the critical time of development when feed is plentiful but lacking energy. Also available with 16% and 20% urea.**

### NUTRIENT ANALYSIS

Protein	56.458%	Sodium	3.725%	Vitamin A	33500 iu/kg
RDP	51.310%	Salt	9.973%	Vitamin D3	4000 iu/kg
UDP	5.147%	Potassium	0.761%	Vitamin E	125 iu/kg
Equiv CP	44.166%	Magnesium	1.114%	Iodine	3.75 mg/kg
Crude Fibre	4.375%	Sulphur	4.134%	Manganese	300 mg/kg
ME RUM MJ	3.895 mj/kg	Urea	8.333%	Zinc	300 mg/kg
Calcium	3.979%	Phosphorous	0.354%	Selenium	0.75 mg/kg

**WARNING: UREA is toxic, keep the product dry and ensure the container drains well. Do not feed to dogs, horses or other equines as may be fatal.**

**The benefits of The Brew include improved liveweight gains, increase productivity and lowering the incidence of weaner death. The Brew can be easily applied to most weaner programs.**

**Urea:** Can be toxic to stock if the rumen is not adequately adjusted, ensure stock can always gain access to the Brew.

**Sulphate of Ammonia:** Supplies sulphur to the bacteria in the rumen and also helps to reduce intake.

**Cottonseed Meal:** Is a very high bypass protein source. (Approx 42%)

**Salt:** Attracts stock, initially salt should be supplied freely for approx 10 days before commencing feeding of the Brew.

**Lime:** Used for calcium delivery.

**Zeolite:** Holds up to 4 times its weight in water which helps to hold the ammonia and release it slowly over time, ensuring the urea works effectively.

#### Directions:

- Salt should be fed in trough or drum for 8 days prior to introducing The Brew 8%.
- Ensure The Brew is kept in a clean dry half drum or trough. Urea may become toxic if wet.
- During periods of wet weather if the product is not well covered remove The Brew and place out salt until it is once again safe to reintroduce. Always ensure an adequate supply of roughage is available for stock.

#### Intake:

- 100 to 150 grams per head per day for cattle and 10 to 15 grams per head per day for sheep approx.
- For Sheep; feed salt for 8 days prior to The Brew 8%, as consumption increases then use The Brew 16%.
- For Cattle; feed salt for 8 days prior to The Brew 8%, as consumption increases then use The Brew 16%, then The Brew 20%.

## Available in a 25kg bag

**Note 1:** Urea is not a protein, rather it is a simple nitrogen compound. Urea may be harmful when supplemented incorrectly. Do not feed to starving animals or in poor condition. Ensure there is sufficient forage available when feeding this product. Be wary when product is wet. Store below 30° C (Room temperature).

**Note 2:** Soil fertility and water quality have substantial effects on stock intake in this free choice grazing management program.

The above feeding rates are a guide only. Variances due to pasture quality, climate, workload and age of the animal will give different results. This product is suitable for ruminants only and not recommended for other animals. Please use the feeding table as a guide only and ensure feeding guide is introduced gradually to all animals, as unobserved diet changes can be detrimental to final results or the animals health.

**[www.furneysstockfeeds.com.au](http://www.furneysstockfeeds.com.au)**  
**Phone (02) 6841 3800 Fax (02) 6881 8717**  
**Email to [sales@furneysstockfeeds.com.au](mailto:sales@furneysstockfeeds.com.au)**