

The Brew Dry Season

26% Urea Dry Lick

The Brew Dry Season lick is high in crude protein equivalent at 98.0% obtained from urea as well as ammonia Sulphate. These non-protein nitrogen sources help cattle maintain any benefits associated with dry standing pasture or pulled edible scrub. Phosphorous is included to assist breeders during dry times. It is well known that some districts have depleted phosphorous in their pasture and that calving percentages increase significantly when a full profile of macro and micro trace elements and minerals are present.

NUTRIENT ANALYSIS

Protein	98.0%	Salt	25.0%	Cobalt	30 mg/kg
Equiv CP	90.5%	Calcium	3.5%	Iodine	30 mg/kg
Crude Fibre	10.0%	Phosphorous	3.5%	Zinc	500 mg/kg
Crude Fat	0.1%	Fluorine	0.1%	Selenium	2.5 mg/kg
Urea	26.0%	Sulphur	4.5%	Molasses	10.0%
Sulphate of Ammonia	15.0%	Copper	300 mg/kg		

WARNING: UREA is toxic, keep the product dry and ensure the container drains well. Do not feed to dogs, horses or other equines as may be fatal.

Ingredients:

Urea, Cottonseed meal, Coconut meal, Ammonium Sulphate, Limestone, Di Calcium Phosphate, Salt, Furneys ruminants vitamin and mineral premix and Liquid Molasses.

**Available
in
25kg bag**

Directions:

- Salt should be fed in trough or drum for 8 days prior to introducing The Brew Dry Season.
- Ensure The Brew Dry Season is kept in a clean dry half drum or trough. Urea may become toxic if wet.
- During periods of wet weather if the product is not well covered remove The Brew Dry Season and place out salt until it is once again safe to reintroduce.
- Always ensure an adequate supply of roughage is available for stock.

Intake: The Brew Dry Season should be approx 100–150gm per head per day for cattle.

For Cattle: Feed out Salt for 8 days prior to using The Brew Dry Season.

Note 1: Urea is not a protein, rather it is a simple nitrogen compound. Urea may be harmful when supplemented incorrectly. Do not feed to starving animals or in poor condition. Ensure there is sufficient forage available when feeding this product. Be wary when product is wet. Store below 30° C (Room temperature).

Note 2: Soil fertility and water quality have substantial effects on stock intake in this free choice grazing management program.

The above feeding rates are a guide only. Variances due to pasture quality, climate, workload and age of the animal will give different results. This product is suitable for ruminants only and not recommended for other animals. Please use the feeding table as a guide only and ensure feeding guide is introduced gradually to all animals, as unobserved diet changes can be detrimental to final results or the animals health.

www.furneysstockfeeds.com.au
Phone (02) 6841 3800 Fax (02) 6881 8717
Email to sales@furneysstockfeeds.com.au