

The Brew Phos

4% Phosphorous Dry Lick

The Brew Phos is a comparatively low protein lick with the view of supplementing sheep of all ages and cattle at the introduction phase. The relatively high level of phosphorous assists breeding ewes and cows with successful joinings and subsequent delivery of healthy offspring. The Brew Phos has found a niche where Mulga is found and used for fodder as Mulga is known to be low in Phosphorous.

NUTRIENT ANALYSIS			
Protein	30.0%	Calcium	11.0%
Equiv CP	25.9%	Fluorine	0.1%
Crude Fibre	10.0%	Copper	300 mg/kg
Crude Fat	0.1%	Cobalt	30 mg/kg
Phosphorous	4.0%	Iodine	30 mg/kg
Urea	9.0%	Zinc	500 mg/kg
Salt	35.0%	Selenium	2.5 mg/kg

WARNING: UREA is toxic, keep the product dry and ensure the container drains well. Do not feed to dogs, horses or other equines as may be fatal.

Ingredients:

Urea, Cottonseed meal, Coconut meal, Ammonium Sulphate, Limestone, Di Calcium Phosphate, Salt, Vegetable oil and Furneys ruminants vitamin and mineral premix.

Directions:

- Salt should be fed in trough or drum for 8 days prior to introducing The Brew Dry Season.
- Ensure The Brew Dry Season is kept in a clean dry half drum or trough. Urea may become toxic if wet.
- During periods of wet weather if the product is not well covered remove The Brew Dry Season and place out salt until it is once again safe to reintroduce.
- Always ensure an adequate supply of roughage is available for stock.

**Available
in
25kg bag**

Intake: The Brew Phos should be approx 100—150gm per head per day for cattle and 10—15gm per head per day for sheep.

For Sheep: Feed out Salt for 8 days prior to using The Brew Phos.

For Cattle: Feed out Salt for 8 days prior to using The Brew Phos.

Note 1: Urea is not a protein, rather it is a simple nitrogen compound. Urea may be harmful when supplemented incorrectly. Do not feed to starving animals or in poor condition. Ensure there is sufficient forage available when feeding this product. Be wary when product is wet. Store below 30° C (Room temperature).

Note 2: Soil fertility and water quality have substantial effects on stock intake in this free choice grazing management program.

The above feeding rates are a guide only. Variances due to pasture quality, climate, workload and age of the animal will give different results. This product is suitable for ruminants only and not recommended for other animals. Please use the feeding table as a guide only and ensure feeding guide is introduced gradually to all animals, as unobserved diet changes can be detrimental to final results or the animals health.

www.furneysstockfeeds.com.au
Phone (02) 6841 3800 Fax (02) 6881 8717
Email to sales@furneysstockfeeds.com.au