

# GREEN FEED BLOCK

AVAILABLE IN 20KG HARD PRESSED BLOCKS



## NUTRITIONAL ANALYSIS

TERIC OIL / ETHOXYLATE ALCOHOL (surfactant - anti foaming agent)	8.3%
MOLASSES	16.0%
PHOSPHOROUS	5.2%
DI CALCIUM PHOSPHATE	5.0%
CALCIUM	8.8%
CALCIUM CARBONATE	4.4%
MAGNESIUM	4.2%
MAGNESIUM SULPHATE	2.1%
MAGNESIUM CARBONATE	2.1%
CRUDE PROTEIN	2.5%
CRUDE FAT	0.5%
CRUDE FIBRE	6.5%
SALT	27%
IRON SULPHATE	0.3%
SULPHUR	1.0%
POTASSIUM IODIDE	0.2%
MANGANESE SULHATE	0.005%
ZINC OXIDE	0.015%
COBALT SULPHATE	0.015%
FLORINE	0.020%

## INGREDIENTS

Molasses, Poloxalene, Teric Oil / Ethoxylate Alcohol, Rock Phosphate, Lime Wheat, Sorghum, Barley, Magnesium Sulphate, Magnesium Carbonate, Salt, Iron Sulphate, Sulphur, Potassium Iodine, Maganese Sulphate, Zinc Oxide, Cobalt Sulphate



## WARNING

This product is suitable for sheep, cattle and goats only and is not recommended for other animals. The above feeding rates are a guide only. Variances due to pasture quality, climate, workload and age of the animal will give different results. Please use the feeding table as a guide only and ensure feeding is introduced gradually to all animals, as unobserved diet changes can be detrimental to final results or the animals health. This product does not contain UREA. This product does not contain restricted animal material. Always ensure clean drinking water is available.

Furneys Green Feed (bloat) Block is designed to assist with bloat, swollen abdomen and acute tympany brought on from fresh green legume pastures. The blocks should be put out before the risk appears and sufficient blocks distributed to ensure that even shy feeders have access to the blocks. Furneys Green Feed (bloat) Blocks are a softer block to allow maximum consumption. This product uses an anti foaming agent in the form of Teric Oil (Ethoxylate Alcohol - Surfactant) which settles on top of the stomach and reduces the production of foam that builds up and causes bloat.



## FEEDING GUIDE

Furneys Green Feed (bloat) Blocks are recommended to be fed based on 1 block per number of head, refer to feeding table below. Feed prior to going into risky pasture.

### \*\*\*BLOAT CONTROL RECOMMENDATION\*\*\*

**Take stock off dangerous pasture and put in small holding paddock overnight.**

**Every morning put out low grade hay or straw to fill gut.**

**When controlled grazing, place bloat oil on water troughs. Place as many blocks as deemed sensible.**

## FEEDING TABLE

ANIMAL TYPE	NUMBER OF HEAD	NUMBER OF DAYS
CATTLE	10	28
SHEEP / GOATS	50	28



**FOLLOW US**  
to keep up with our latest news, specials & competitions!



## DUBBO

115 Erskine Street  
(02) 6841 3800

## ORANGE

245 McLachlan Street  
(02) 6362 2896

## RUTHERFORD

78 Racecourse Road  
(02) 4932 4455

## TAMWORTH

7 Wallamore Road  
(02) 6762 1363

## TOOWOOMBA

526-528 Boundary Road  
(07) 4616 1270