Feeding the country's livestock for over 101 years

LICK OR LOT + RUMENSIN

AVAILABLE IN 25KG BAGS





Added RUMENSIN To Increase Energy & Reduce Bloat

RUMENSIN™ increases the efficiency of the energy available from all feed consumed by the animal, reduces the incidence of feedlot bloat and enhances productivity by increasing the rate of weight gain and improving feed conversion efficiency of the animal.



Feeding Guide

FEEDING WITH DRY ROUGHAGE

Furneys Lick or Lot + Rumensin can be used as a supplement in lick feeders or troughing at 250g per head per day for weaner type cattle with cereal grain (cracked barley preferably.) Starting – Pour limited amount into trough, make sure stock are introduced to this until you are satisfied all stock are licking the supplement. Then add grain mix. Ensure cattle have access to pasture at all times.

FEEDING AS A FEEDLOT ADDITIVE

Furneys Lick or Lot + Rumensin can be fed as a feedlot additive. We recommend feeding Furneys Lick or Lot + Rumensin including 120kg of lick with 750kg of grain (barley cracked). Have either plenty of dry grass in the paddock or include hay.



DAILY INTAKE	
ANIMAL TYPE	APPROXIMATE FEEDING RATE
Cattle	250 grams per head per day.
Sheep	25 to 30 grams per headper day.
WITHHOLDING PERIOD	
Cattle	NIL Withholding Period
Sheep	1 Day Standard OR 7 Day Export Slaughter Withholding Period

Note: Protein availability: Cotton Seed Meal is the most effective bypass protein meal. If not in production, Furneys use Canola Meal or Soyabean Meal.










Ingredients

Cotton Seed Meal and/or Canola Meal and/or Copra Meal and/or Soyabean Meal, Millrun, Acid Buf, Sodium Bentonite, Lime, Salt, Urea, Rumensin 10% Premix and Furneys Cattle Premix.



FOLLOW US

to keep up with our latest news, specials & competitions!





WARNING DO NOT FEED TO DOGS, HORSES OR OTHER EQUINES AS IT MAY BE FATAL. RUMENSIN IS NOT REGISTERED FOR GOATS OR DEER.

This product is suitable for sheep and cattle only and not recommended for other animals. The above feeding rates are a guide only. Variances due to pasture quality, climate, workload and age of the animal will give different results. Please use the feeding table as a guide only and ensure feeding is introduced gradually to all animals, as unobserved diet changes can be detrimental to final results or the animals health. This product contains Urea, keep the product dry and ensure it drains well. This product does not contain restricted animal material. Always ensure clean drinking water is available.

DUBBO

115 Erskine Street (02) 6841 3800

ORANGE

245 McLachlan Street (02) 6362 2896

RUTHERFORD

78 Racecourse Road (02) 4932 4455

TAMWORTH

7 Wallamore Road (02) 6762 1363

TOOWOOMBA

526-528 Boundary Road (07) 4616 1270